arXiv:astro-ph/0408464v1 25 Aug 2004

Is the Radion a Chameleon?

Ph. Brax

Service de Physique Théorique, CEA–Saclay, Gif/Yvette cedex, France F–91191

C. van de Bruck

Astro–Particle Theory and Cosmology Group, Department of Applied Mathematics
University of Sheffield, Sheffield S3 7RH, UK

A.–C. Davis

DAMTP, Centre for Mathematical Sciences, Cambridge University
Wilberforce Road, Cambridge CB3 0WA, UK

Abstract. The chameleon mechanism is a way to give an effective mass to a light scalar field via field self-interaction and interaction with matter. We study this mechanism in models in which the couplings are field dependent and find that the properties are very different from the case of constant couplings. The consequences of a runaway potential for the radion field in brane world scenarios and whether the radion can play the rôle of dark energy is investigated. The cosmological evolution during the inflationary epoch, the radiation and the matter dominated epochs are discussed as is the compatibility of the radion field with local tests of gravity.

1. Introduction

Current cosmological observations suggest that the expansion of the universe is accelerating instead of slowing down (see e.g. [1] for recent results). The type of matter responsible for the accelerated expansion has been dubbed *dark energy*. Not much is known about the nature this new form of energy but different candidates have been suggested such as a cosmological constant or a slowly rolling scalar field (also called quintessence–field or cosmon–field) (see [2] and [3] for early work and e.g. [4, 5] for recent reviews). A challenge for particle physics is to identify candidates with the right properties. For example, the coupling of any new matter type to ordinary matter cannot be too large, otherwise dark energy would have been detected in the form of an additional force acting on bodies in laboratory experiments.

If current observations are confirmed by future experiments, it is highly probable that an explanation for dark energy will involve new physics beyond the standard models of particle physics and cosmology. One possible type of new physics beyond the standard model is the idea of extra dimensions. Among the models with higher dimensions, brane worlds have attracted a lot of attention recently. For recent reviews see [6]–[10]. It has been speculated that the current accelerated expansion is due to the effects of extra spatial dimensions [11, 12, 13, 14, 15, 16]. Such models can be tested (or constrained) with the observation of the CMB anisotropies, see e.g. [17] and [18]. Higher dimensional theories provide a plethora of scalar fields which usually couple to matter. String theory, for example, predicts the existence of the dilaton field, which couples universally to all matter forms. Another example is the brane world theory, according to which our universe is a hypersurface embedded in a higher dimensional space. There, scalar fields in the bulk and/or the size of the extra dimensional spacetime manifest themselves as scalar fields in the low–energy description of the system. Usually, all these fields couple to all matter forms confined on the brane. It is so far not known if these fields can play the rôle of dark energy and, at the same time, remain hidden from experiments on earth. Several proposals have been made in the past, in which the quintessence field couples to baryons and/or dark matter (see e.g. [19]–[22] and references therein). Cosmological considerations, such as nucleosynthesis, the anisotropy spectrum of the cosmic microwave background radiation, structure formation, etc., constrain the coupling of the quintessence field to matter (baryonic or non–baryonic).

In this paper we investigate a system of two branes and bulk scalar field as a specific setup of a higher–dimensional theory. We discuss under which conditions the radion, i.e. the interbrane distance, can be a candidate for dark energy and if, at the same time, the theory is compatible with local experiments. We are going to assume that a potential for the radion field has been generated, for example by non–perturbative effects. Therefore, in our model the dynamics of the radion is governed by scalar field self-interactions and interaction with surrounding matter. Such fields, in which the effective potential is matter (environment) dependent, have recently been dubbed *chameleon fields* and arise usually in scalar–tensor theories with a potential for the scalar field [23]. We show that for scalar–tensor theories with an essentially linear coupling function, one can evade local tests thanks to the existence of a thin–shell regime: the chameleon field is basically constant inside a big, spherical body and varies only inside a thin–shell close to the outer edge of the body. This implies that the fifth force outside the body is a small correction to Newton's law.

There are two regimes for the radion in the brane model we consider. At small distances (i.e. distances smaller than the bulk curvature scale), the radion possesses the thin–shell property while at large distance (distances larger than the bulk curvature scale) no thin-shell regime exists. The thin–shell case is expounded in [23, 24]. Here we focus on the case of thick–shells at large distances between the branes.

The paper is organized as follows: In Section 2 we describe briefly the chameleon properties. In particular, thin and thick shells are discussed. In Section 3 we focus on the radion and show that both possibilities are realised at either small or large distances. In Section 4 we investigate first the inverse power–law potential in detail. After discussing the cosmological evolution, we elaborate on local experiments and constraints. These constraints are stringent. We then generalise these results to a class of potentials where, at large distances between the branes, the radion can both be a quintessence field and comply with gravity experiments. In Section 5 we discuss a non–standard potential and the consequences for the evolution of the radion. We present our conclusion in Section 6.

2. Chameleon Fields

In this Section, we discuss the properties of chameleon fields. Because these fields appear naturally in scalar–tensor theories of gravity in which the scalar field has a potential, we begin with a discussion of these theories.

2.1. Scalar–tensor theories

We consider theories where a scalar field ϕ couples both to gravity and matter, generating a potential fifth force. The Lagrangian of such scalar–tensor theories reads

$$S = \frac{1}{2\kappa_4^2}\int d^4x\sqrt{-g}(R - (\partial\phi)^2 - 2\kappa_4^2 V(\phi)) \tag{1}$$

Notice that ϕ is dimension–less. Redefining $\phi \to \kappa_4^{-1}\phi$ leads to a dimension–one field. Here, $\kappa_4 = 1/m_{\rm Pl}$, where m_{pl} is the reduced Planck mass. Matter couples to both gravity and the scalar field according to

$$S_m(\psi, A^2(\phi)g_{\mu\nu}), \tag{2}$$

where ψ is a matter field and A is an arbitrary function of ϕ. The Klein–Gordon equation for the scalar field involves a coupling to the trace of the matter energy–momentum tensor T

$$D^\mu D_\mu\phi = \kappa_4^2\frac{\partial V(\phi)}{\partial\phi} - \kappa_4^2\alpha_\phi T, \tag{3}$$

where

$$\alpha_\phi = \frac{\partial \ln A}{\partial\phi}. \tag{4}$$

The energy–momentum tensor in the Einstein frame fulfills

$$D_\mu T^\mu_\nu = \alpha_\phi\,(D_\nu\phi)\,T. \tag{5}$$

In a Friedmann–Robertson–Walker Universe these equation simplify to

$$\ddot\phi + 3H\dot\phi = -\kappa_4^2\frac{\partial V(\phi)}{\partial\phi} - \kappa_4^2(1-3w_m)\rho\frac{\partial A}{\partial\phi}, \tag{6}$$

together with

$$\dot\rho + 3H\rho = \alpha_\phi\dot\phi\rho. \tag{7}$$

Redefining

$$\rho = A(\phi)\rho_m, \tag{8}$$

one can easily verify that

$$\rho_m = \frac{\rho_0}{a^{3(1+w_m)}} \tag{9}$$

is the usual energy density.

The Klein–Gordon equation can be written in terms of an effective potential

$$\ddot{\phi} + 3H\dot{\phi} = -\kappa_4^2 \frac{\partial V_{\rm eff}(\phi)}{\partial \phi}, \tag{10}$$

where

$$V_{\rm eff}(\phi) = V(\phi) + \rho_m (1 - 3w_m) A(\phi). \tag{11}$$

The effective potential depends on the environment through ρ_m. In particular, we will assume that $V(\phi)$ is a runaway potential and for the models we consider $A(\phi)$ increases with ϕ. The potential has a minimum for

$$\frac{V'(\phi_{\rm min})}{A'(\phi_{\rm min})} = -\rho_m (1 - 3w_m). \tag{12}$$

Such a field has been called a chameleon field. The field, when sitting at the minimum, becomes massive with mass $(\kappa_4^2 V''_{\rm eff})$

$$m^2|_{\phi_{min}} = -\kappa_4^2 \alpha_\phi (1 - 3w_m) A \frac{d\rho}{d\phi(\rho)}|_{\phi_{min}}, \tag{13}$$

evaluated at the minimum. In this equation, we have made use of the fact that, at the minimum, ϕ is a function of ρ, as seen from equation (12).

Dimensionally the mass is proportional to $\kappa_4^2 \rho_m \approx H^2$ on cosmological scales, i.e. the field at the minimum has a very small mass. Naively, one would expect that in that case solar system experiments would lead to very strong constraints on the models. However, because the effective mass depends on the energy density of the environment, this is not necessarily true.

2.2. Solar Tests

In the following, we impose constraints springing from local experiments, i.e. solar system experiments. The field ϕ acts on all types of matter and, in the Einstein frame, there is a new force associated with the scalar field. Therefore, massive particles no longer follow geodesics with respect to the Einstein frame metric. The equations of motion in the Einstein frame are

$$\ddot{x}^\mu + \Gamma^\mu_{\nu\lambda} \dot{x}^\nu \dot{x}^\lambda + \alpha_\phi \frac{\partial \phi}{\partial x_\mu} = 0 \tag{14}$$

In this equation, x^a represents the four–vector describing the worldline of a test particle moving in the metric $g_{\mu\nu}$, $\Gamma^\mu_{\nu\lambda}$ represents the Christoffel symbols for the metric $g_{\mu\nu}$. The last term is a new contribution, representing the force from the scalar field, which is given by

$$F_\phi = -m\alpha_\phi \frac{\partial \phi}{\partial x_\mu}, \tag{15}$$

where m is the mass of the test particle. The force F_ϕ cannot be too large, otherwise experiments would have already detected it. What is the profile $\phi(r)$ for an object like the earth? Modelling the earth as a spherical object with constant density $\rho_\bullet$, the equation for ϕ (i.e. the Klein–Gordon equation), reads for $r \leq R_\bullet$

$$\phi'' + \frac{2}{r}\phi' = \kappa_4^2 \frac{\partial V_{\text{eff}}}{\partial \phi}, \tag{16}$$

whereas for $r > R_\bullet$ we assume that the field is close to its minimum ϕ_∞ determined cosmologically (we assume that space is empty outside the body)

$$\phi'' + \frac{2}{r}\phi' = m_\infty^2(\phi - \phi_\infty) \tag{17}$$

For the function $A(\phi)$, we will focus on the form

$$A = 1 + \beta \frac{\phi^{m+1}}{m+1}, \tag{18}$$

In this paper, we will consider the case in which β is of order one. As we will show when $m = 0$ a thin–shell regime exists, whereas for $m = 1$ there are no thin shells even for massive bodies. The case $m \geq 1$ is pathological as it leads to no thin-shell and an infinite repulsive force in the core of the spherical body. Putting $\phi = r^{-1}\psi$ we get

$$\psi'' = \beta\kappa_4^2\rho_\bullet \frac{\psi^m}{r^{m-1}}. \tag{19}$$

Let us discuss the case $m = 0$ first.

2.3. The thin–shell regime

We will solve the equations of motion inside and outside the spherical body. We will assume that the minimum outside the body ϕ_∞ is greater that the minimum inside ϕ_b. This is the case for simple potentials like an inverse power law.

First of all, we impose that $\phi' = 0$ at $r = 0$. Interpreting the equations of motion as classical mechanics in the effective potential $-V_{eff}(\phi)$, this corresponds to no velocity at the origin. Moreover we fix the value of the field there to be near the minimum ϕ_b of the effective potential inside the body. The solution can be approximated in three different regions. First for $r \leq R_s$ the field is constant $\phi = \phi_b$. In a shell $R_s \leq r \leq R_\bullet$ the field is increasing in order to match the value $\phi_\infty > \phi_b$. It reads

$$\phi = \phi_b - \frac{3\beta}{8\pi}\frac{\kappa_4^2 M_s}{R_s} + \frac{\beta\kappa_4^2\rho_\bullet}{6}r^2 + \frac{\beta\kappa_4^2}{4\pi}\frac{M_s}{r}. \tag{20}$$

Outside the body for $r > R_\bullet$ the field behaves like a Yukawa potential

$$\phi = \phi_\infty + \frac{\beta\kappa_4^2}{4\pi}\frac{M_s - M_\bullet}{r}e^{-\kappa_4 m_\infty (r - R_\bullet)}. \tag{21}$$

In the vicinity of the body and using the fact that $m_\infty \approx H_0$ we can neglect the exponential profile. Hence the outside the body the force is a correction to Newton's law. Matching at $R_\bullet$ leads to

$$\frac{R_\bullet - R_s}{R_\bullet} = \frac{1}{6\beta}\frac{\phi_\infty - \phi_b}{\Phi_0}, \tag{22}$$

where $\Phi_0 = \kappa_4^2 M_\bullet / 8\pi R_\bullet$ is the gravitational potential on the surface of the body. Because ϕ can be viewed as a potential for a new, fifth force (see eq. (15)), the above equation says that if the ratio between the difference of the field at infinity and inside the body and the gravitational potential is small, R_s is of order $R_\bullet$. This regime has been called the *thin–shell regime*. It is realised when

$$(\phi_\infty - \phi_b) \ll \beta\Phi_0. \tag{23}$$

The new force leads to a modification to Newton's law outside the body

$$F = (1+\theta)F_{Newton}, \tag{24}$$

where $F_{Newton} = -mM_\bullet G_N/r^2$ and

$$\theta = \frac{9G_N(\phi_\infty - \phi_b)}{R_\bullet^2}. \tag{25}$$

which is suppressed

$$|\theta| \ll \frac{\rho_\bullet}{m_{\rm pl}^4} \ll 1. \tag{26}$$

Despite being extremely light, the chameleon field does not generate a measurable fifth force. This results springs from the smallness of $R_\bullet - R_s$, i.e. the thin–shell regime.

To conclude, in the case $m = 0$ we have found that, like in the original chameleon model, the thin–shell regime may exist. We will see that in the case $m = 1$, the thin–shell regime does not exist.

2.4. The thick–shell regime

Let us repeat the same analysis for $m = 1$. In the range $R_s \le r \le R_\bullet$ we find that ϕ is a combination of $\frac{\sinh(r\sqrt{\kappa_4^2\rho_\bullet})}{r}$ and $\frac{\cosh(r\sqrt{\kappa_4^2\rho_\bullet})}{r}$. Matching at R_s leads to

$$R_s = \sqrt{\frac{2}{\kappa_4^2\rho_\bullet}}, \tag{27}$$

implying that $R_s \gg R_\bullet$. Thus, there is no thin–shell regime, even inside big, heavy bodies.

Demanding that the first derivative vanishes at the origin, the solution inside the body can be found to be

$$\phi_I = \tilde{A}\frac{\sinh(r/R_r)}{r}, \tag{28}$$

where $R_r^2 = m_{pl}^2/\beta\rho_\bullet$. Outside the body, the solution reads

$$\phi_E = \mathcal{C} + \frac{\mathcal{D}}{r}. \tag{29}$$

Matching both solutions and their first derivative at $r = R_\bullet$ and noting that $R_\bullet \ll R_r$ and $\phi \to \phi_\infty$ as $r \to \infty$ one obtains

$$\tilde{A} = -3\mathcal{D}\frac{R_r^3}{R_\bullet^3}, \tag{30}$$

$$\mathcal{C} = \phi_\infty, \tag{31}$$

$$\mathcal{D} = \frac{\phi_\infty R_\bullet}{1 + 3\frac{R_r^2}{R_\bullet^2}}, \tag{32}$$

where ϕ_∞ is the cosmological value of the field now, i.e. the value of the field far away from the body.

Thus, away from the body, the profile is given by

$$\phi(r) = \phi_\infty - \frac{\beta\phi_\infty}{4\pi}\frac{M_\bullet}{m_{pl}^2}\frac{1}{r}, \tag{33}$$

leading to a force

$$F_\phi = -\frac{\beta^2\phi_\infty^2}{4\pi}\frac{mM_\bullet}{m_{pl}^2 r^2}, \tag{34}$$

which is nothing but a correction to Newton's law with

$$\theta = 2\beta^2\phi_\infty^2. \tag{35}$$

Thus, in the case with $m = 1$, the effective coupling between scalar field and matter is not specified by β alone, *it is also determined by the cosmological value of the field itself.* Experiments constrain $\theta \leq 10^{-5}$, see [25] and [26]. This implies that the value of the field at infinity, determined by the cosmological dynamics, must be extremely small.

2.5. The thick–shell catastrophe

We will briefly discuss the case $m > 1$. The solution inside the body reads

$$\phi = F_0 r^{2/(1-m)} \tag{36}$$

where $F_0^{m-1} = \frac{2(3-m)}{(1-m)^2\beta\kappa_4^2\rho_\bullet}$. This solution is only valid when m is even, the $(m-1)$–th root taking alternative signs. Notice first that this solution cannot match a constant between R_s and $R_\bullet$. Hence there is no thin–shell and the above solution is valid all the way to the origin. At the origin the field diverges and the force behaves like

$$\frac{F}{m} = O\left(\frac{1}{r}\right) \tag{37}$$

It is repulsive and infinite at the origin implying that the compact spherical object would not exist.

In conclusion, only $m = 0, 1$ are sustainable cases. Whereas the case $m = 0$ allows for a thin–shell regime, the case $m = 1$ does not have this propery. The case $m > 1$ is pathological and does not lead to physical acceptable results.

3. The Radion

Let us now turn our attention to a scalar–tensor theory motivated from the brane world idea. In what follows, we will in particular focus on the radion field, i.e. the interbrane distance. Let us first concentrate on two-brane models with a bulk scalar field (for details, see e.g. see [27], [28] and [29]). At low energy the model is described by a tensor–scalar field theory with two massless scalar fields, $\mathcal{R}$ and $\mathcal{S}$, related to the scale factor in the bulk $a(z)$

$$\tanh\frac{\mathcal{R}}{\sqrt{6}} = \frac{a^{1+2\delta^2}(z_2)}{a^{1+2\delta^2}(z_1)} \tag{38}$$

where $z_{1,2}$ are the brane coordinates and

$$\mathcal{S} = \frac{1}{2}\ln(a^{2+4\delta^2}(z_2) - a^{2+4\delta^2}(z_1)) \tag{39}$$

For $\delta = 0$, the second field $\mathcal{S}$ decouples and we obtain the Randall-Sundrum model [30] with a single field

$$d = -l\ln\left(\tanh\frac{\mathcal{R}}{\sqrt{6}}\right) \tag{40}$$

where d is the interbrane distance, i.e. the radion, and $l = 1/k$ measures the curvature scale of the bulk space– time. However, even if $\delta \neq 0$ we will still call $\mathcal{R}$ the radion field.

In the bulk, the scale factor behaves like

$$a(z) = (1 - 4k\delta^2 z)^{1/4\delta^2} \tag{41}$$

showing that there is a would–be singularity when $a = 0$ beyond the second brane. When coupled to matter, the cosmological evolution drives $\mathcal{R}$ to zero, i.e. the second brane hits the would–be singularity [29, 27].

This situation can be avoided by introducing a potential $V(r)$ shielding $\mathcal{R} = 0$. Such a run–away potential can be chosen to be like

$$V(\mathcal{R}) = \frac{\Lambda^4}{(\mathcal{R} - \mathcal{R}_*)^\alpha}. \tag{42}$$

where $\mathcal{R} \geq \mathcal{R}_*$ is the allowed range. As we assume that $\mathcal{S}$ is still massless, solar system experiments imply that $\delta \ll 1$, i.e. the Randall–Sundrum limit. In that case the coupling of the field $\mathcal{R}$ to matter is well specified by the function [29]

$$A(\mathcal{R}) = \cosh\left(\frac{\mathcal{R}}{\sqrt{6}}\right). \tag{43}$$

In the Randall-Sundrum case, there are two relevant regimes. When $\mathcal{R} \gg 1$, the distance between the branes is small $d \ll l$ (i.e. the distance is smaller than the bulk curvature length scale). Changing $\mathcal{R} \to \tilde{\mathcal{R}} = \mathcal{R} - \mathcal{R}_*$, the coupling becomes then

$$A(\mathcal{R}) = \frac{1}{2}e^{\frac{\mathcal{R}+\mathcal{R}_*}{\sqrt{6}}} \tag{44}$$

In that case, we retrieve the original example of a chameleon field corresponding for $\mathcal{R}_* \gg 1$ and $\tilde{\mathcal{R}} \ll 1$ to a linear coupling with $m = 0$. Hence the radion possesses then a thin–shell. This situation has been studied at length in [23]. Now there is another interesting regime when $\mathcal{R} \ll 1$ and $d \gg l$ (i.e. interbrane distances larger than the bulk curvature scale), where we can approximate

$$A \approx 1 + \frac{\mathcal{R}^2}{12}, \tag{45}$$

and we have taken $\mathcal{R}_* = 0$. It corresponds to

$$m = 1 \quad \text{and} \quad \beta = \frac{1}{6} \tag{46}$$

in equation (18).

Thus, if $\mathcal{R}$ is small enough with a runaway potential of the form (42), we end up with a theory in which the thin–shell regime does not exist even for heavy bodies. Cosmology has to ensure that the field is small today and fullfills nucleosynthesis constraints. We will therefore now discuss the cosmology of the radion field with a potential energy of the form (42) and therefore get an estimate of ϕ_∞, which determines the coupling to matter locally (see eq. (35)).

4. Quintessential Potentials for the Radion

In this section, we will switch notation and will denote the radion field as ϕ. We will discuss the different cosmological epochs separately. Before imposing the local constraints, however, we will discuss the cosmological evolution in general.

4.1. The inflationary era

Let us consider first an inflationary era in order to find the initial conditions for the radion field in the radiation dominated epoch. For the inflationary model, we choose a chaotic inflationary potential of the form

$$V(\sigma) = \frac{1}{2} m^2 A^2 \sigma^2. \tag{47}$$

where σ has dimension one. We assume that the inflaton field σ is confined on the positive tension brane. Then, in the Einstein frame, its mass depends on the moduli fields through a factor of A. The Klein–Gordon equation for the field ϕ reads

$$\ddot{\phi} + 3H\dot{\phi} + \kappa_4^2 \frac{\partial V(\phi,\sigma)}{\partial \phi} = -\kappa_4^2 T \alpha_\phi \tag{48}$$

where $T = -\dot{\sigma}^2 - 2m^2 A^2 \sigma^2$ is the trace of the energy momentum tensor of σ. Thus, the dynamics of the field ϕ are governed by an effective potential

$$V_{\text{eff}} = \frac{\Lambda^4}{\phi^\alpha} + \frac{3}{2}(1 + \frac{\beta}{2}\phi^2) m^2 \sigma^2, \tag{49}$$

where we have neglected the kinetic energy of σ as the field is assumed to slow–roll. During inflation, $\sigma = \sigma_{\text{inf}}$ is almost constant. The minimum of the effective potential is given by

$$\phi_0 = \left(2 \frac{\alpha \Lambda^4}{3\beta m^2 \sigma_{inf}^2} \right)^{1/(\alpha+2)} \tag{50}$$

and the effective mass of the ϕ field at the minimum is

$$m_{\text{eff}}^2 = 3\beta(\alpha+2) H^2, \tag{51}$$

where $H^2 = m^2 \sigma_{\text{inf}}^2 / 2 m_{\text{Pl}}^2$. This large mass guarantees that the effective potential for ϕ is very steep around its minimum and ϕ decays rapidly towards the minimum. Thus, just before the end of inflation, ϕ sits at the minimum of the effective potential. Notice that the value of ϕ is tiny, as Λ is small. We will assume that the field ϕ is a fundamental field and therefore that the inflaton field does only decay into radiation, the standard model particles and dark matter. Thus, during reheating no energy will be released into the field ϕ and therefore during reheating the field remains small. This provides a hint about the initial value of the radion at the beginning of the radiation epoch.

4.2. The radiation– and matter–dominated epoch

Let us consider now the cases of the radiation and matter dominated epochs. The Klein–Gordon for the radion equation reads

$$\ddot{\phi} + 3H\dot{\phi} + \kappa_4^2 \frac{\partial V(\phi)}{\partial \phi} = -\kappa_4^2 \rho_m \phi, \tag{52}$$

that is, the effective potential reads

$$V_{\rm eff} = \frac{\Lambda^4}{\phi^\alpha} + \rho_m \left(1 + \frac{\beta}{2}\phi^2\right). \tag{53}$$

The field value at the minimum is given by

$$\phi_{\rm min} = \left(\frac{\alpha \Lambda^4}{\beta \rho_m}\right)^{1/(\alpha+2)}. \tag{54}$$

Because the field is small, we have that

$$\rho_m \approx \frac{\rho_m^0}{a^3}. \tag{55}$$

Now we can consider two cases during radiation domination: either the fields starts at $\phi(t_i) > \phi_{\rm min}$ or at $\phi(t_i) < \phi_{\rm min}$. Assume that the field value is larger than the minimum. Then we can neglect the potential and the field is governed by

$$\ddot{\phi} + 3H\dot{\phi} \approx -\frac{\rho_m}{m_{pl}^2}\phi. \tag{56}$$

This is the equation of an damped oscillator. Because the mass is much smaller than the Hubble parameter, the field is overdamped and does not move. Only if H drops below the mass the field starts to evolve and moves towards the minimum. Let us consider now the case that $\phi < \phi_{\rm min}$. If the potential energy is initially very large, the field rolls down and becomes kinetically dominated (Ω_i is the initial fraction of energy stored in the radion field) (see also [31])

$$\phi = \sqrt{6\Omega_i}(1 - \frac{a_{\rm ini}}{a}) \tag{57}$$

with an equation of state $w = 1$. It stops at

$$\phi_{\rm stop} = \sqrt{6\Omega_i} \tag{58}$$

with an equation of state $w = -1$. This can be seen in figure 3 where a plateau is reached before converging to the attractor (in Planck units)

$$\phi_{\rm rad}(t) = \left(\frac{2\alpha\Lambda^4}{(2\gamma^2 + \gamma)m_{pl}^4}\right)^{1/(2+\alpha)} (m_{pl}t)^\gamma, \tag{59}$$

with $\gamma = 2/(2+\alpha)$. The equation of state is given by

$$w = \frac{(\alpha - 2)\gamma - 1}{(\alpha + 2)\gamma + 1} \tag{60}$$

corresponding to the usual Ratra–Peebles attractor [32].

In the matter dominated era, the minimum of the effective potential is located at

$$\phi_{\rm min}^{\alpha+2}=\frac{\alpha\Lambda^4}{\rho_m} \tag{61}$$

The mass at the minimum (V'') is then

$$m^2=(\alpha+2)\rho_m \tag{62}$$

in Planck units. The Klein–Gordon equation has an attractor

$$\phi_{\rm matt}(t)=\left(\frac{\alpha\Lambda^4}{m_{pl}^4(\gamma^2+\frac{\gamma}{2}+\frac{4}{3}\beta})\right)^{\frac{1}{\alpha+2}}(m_{pl}t)^{\gamma} \tag{63}$$

implying that

$$\frac{\phi_{\rm matt}}{\phi_{\rm min}}=\left(\frac{4}{4\beta+3\gamma^2+3\frac{\gamma}{2}}\right)^{\frac{1}{\alpha+2}} \tag{64}$$

Notice that $\phi_{\rm matt}<\phi_{\rm min}$. The field follows the minimum but does not settle at the minimum (see figure 3). The equation of state reads

$$w=\frac{(\alpha-2)\gamma^2-\gamma-\frac{8}{3}\beta}{(\alpha+2)\gamma^2+\gamma+\frac{8}{3}\beta}, \tag{65}$$

which is smaller than in the radiation dominated era.

We have numerically solved the equations and compared it to the ordinary quintessence model in General Relativity. The results are shown in Figures 1, 2 and 3.



Figure 1. Evolution of the equation of state for the model in General Relativity (dashed line) and in the brane model (solid line) as a function of $1+z$. It can be seen that the equation of state today is nearer -1 in the brane model than in General Relativity. The exponent in the inverse power–law potential is $\alpha=1$, for which $\gamma=2/3$.



Figure 2. Dependence of the equation of state as a function of $\Omega_{\rm m}$. As can be seen, the dependence is weaker in the brane model (solid line) than in General Relativity (dashed line). For this plot, the exponent in the inverse power–law potential is $\alpha = 1$.

In Figure 1 we plot the evolution of the equation of state as a function of $1 + z$ for the case of $\alpha = 1$ and the same initial conditions. We compare the brane theory together with a quintessence model in General Relativity. As can be seen, the evolution of the equation of state around $z = 0$ is more pronounced in General Relativity than in the brane model. Furthermore, the equation of state is smaller in the brane scenario than in General Relativity.



Figure 3. Evolution of the field as a function of $1 + z$ for different initial conditions. The straight dashed line shows the position of the minimum of the effective potential.

In Figure 2 we show the dependence of the equation of state w as a function of $\Omega_{\rm m}$. It can be seen that in General Relativity this dependence is stronger than in the brane model. Finally, in Figure 3 we plot the evolution of the radion field (in Planck units) as a function of $1+z$ for different initial conditions. As can be seen, the field evolves according to the solution (59) in the radiation dominated epoch and according to (63) in the matter dominated epoch.

4.3. Imposing local constraints

The discussion on the cosmological evolution so far did not take into account the constraints imposed by local experiments. In fact, if the radion field acts as dark energy, its evolution is similar to a field with inverse power–law potential in General Relativity. In particular, for $\alpha = \mathcal{O}(1)$, the field value today is of order the Planck mass. This, however, is not compatible with local experiments. Imposing that $\theta \leq 10^{-5}$ leads to $\Lambda \approx 10^{-3}$ eV and $\alpha \leq 10^{-5}$. The potential is extremely flat and peaked sharply around $\phi = 0$. Thus, for the potential considered here the price to be paid is that, at least for the standard potential used here, the parameters have to be extremely fine–tuned, if the radion should play the role of dark energy.

5. The case for a non–standard potential

So far we have focused on a particular inverse power law potential, in which $V(\phi) \to 0$ for $\phi \to \infty$. In the following we will consider another class of potentials leading to an interesting cosmology with no conflict with local experiments. The potential is not a quintessential potential, however, but has the form

$$V(\phi) = M^4 e^{\left(\frac{M}{m_{pl}\phi}\right)^n} \tag{66}$$

where $M = 10^{-3}$ eV in order to trigger acceleration now. It is not a quintessence potential per se as it does not converge to zero at infinity. It should be considered as a very flat potential which may appear from some non-perturbative physics for the radion. As we will now see, such a potential leads to acceleration and drives the field towards very small values ($\phi_\infty \ll 1$).

For the potential above, the minimum of the effective potential is given by

$$\phi_{min}^{n+2} = n \left(\frac{M}{m_{pl}}\right)^n \frac{V(\phi_{min})}{\beta \rho_m}. \tag{67}$$

For $V = O(M^4)$ this leads to

$$\phi_\infty^{n+2} = O\left(\frac{M}{m_{pl}}\right)^n. \tag{68}$$

Notice that $\phi_\infty = O(10^{-10})$ for $n = 1$ implying that solar tests are automatically satisfied.

Let us discuss the cosmology of this model. The mass of the radion at the minimum reads

$$\frac{m^2}{H^2} = 3\beta\Omega_m \left[n(n+2) + n^2(\frac{M}{m_{pl}})^n \frac{1}{\phi^n}\right]. \tag{69}$$

There are two regimes. When $m_{pl}\phi \ll M$, the mass is very large $M \gg H$ implying that the minimum acts as an attractor. When $m_{pl}\phi \gg M$, the potential reads

$$V(\phi) = M^4 + \frac{\Lambda^4}{\phi^n}, \tag{70}$$

where

$$\Lambda^4 = \frac{M^{4+n}}{m_{pl}^n}. \tag{71}$$

Now as long as $V(\phi)$ is subdominant, the constant term is irrelevant and we retrieve the analysis of the power law potential. This implies that in the radiation and matter dominated eras, the radion follows the attractors that we found previously. Only the early Universe is modified.

We have seen that $\phi_{now} \ll 1$ and therefore solar tests are satisfied. What about laboratory tests of the equivalence principle? Let us consider the gravity experiments in a cavity of radius R_*. Inside the vacuum chamber the field is early constant and determined by

$$V''|_{\phi_{cav}} = m_p^2 R_*^{-2}, \tag{72}$$

i.e. the Compton wavelength of the radion is then given by R_*. Moreover the field in the cavity is larger than the field outside the cavity implying that

$$|\frac{\partial\phi}{\partial r}| \leq \frac{\phi_{cav}}{R_*}. \tag{73}$$

Inside the cavity, a gradient of ϕ leads to a new force acting on particles and a breaking of the equivalence principle to the accuracy level

$$\eta = 10^{-4}\beta\frac{\phi_{cav}^2}{R_*}\frac{4\pi m_{pl}^2 R_{earth}^2}{M_{earth}} \tag{74}$$

Imposing that $\eta \leq 10^{-26}$ [26] leads to

$$\phi_{cav} \leq 10^{-13}. \tag{75}$$

which is always satisfied.

To conclude this Section, for a potential of the form (66) the mass scale has to be chosen such that $M \approx 10^{-3}$ eV if the radion is to be a dark energy candidate. This tuning is no more than that required by a cosmological constant.

6. Conclusions

In this paper, the chameleon mechanism has been investigated in the case of field dependent couplings. Such couplings arise naturally in theories with extra dimensions, such as brane world scenarios. In theories with constant couplings a thin–shell regime exists, in which the deviations from General Relativity are suppressed. We have shown, that this is not true in models with field–dependent couplings. In these models the thin–shell regime does not exist (i.e. no suppression of the couplings) and, hence, there are potentially large deviations from General Relativity.

Using the low-energy effective action, we have considered the cosmology of the radion field at large distances on the branes (or low energy). Possible deviations from General Relativity due to the smallness of the radion mass were investigated. Two regimes can be distinguished, leading to the thin and thick shell properties. At short distances between the branes, the radion satisfies the thin-shell property and no deviation can be detected either on earth or in solar system experiments. Satellites experiments would be able to detect large deviations from gravity. For large distances between the branes, the thin-shell property is replaced by a thick–shell regime. In this case deviations from normal gravity are small provided that the cosmological value of the radion is small enough. We have investigated a class of potentials for which the radion (at large distances between the branes) can both act as quintessence and satisfies gravity tests. Unfortunately, future satellite experiments would not notice the presence of the radion as the coupling in that case is too small.

If the branes are close together, the model and its cosmology are similar to the original chameleon model [23, 24], local constraints are satisfied and the radion can also act as a dark energy candidate. In this case, future satellite experiments could detect the predicted deviations from General Relativity.

Acknowledgements: We are grateful to Justin Khoury for discussions. We wish to thank the British Council–Alliance for an exchange grant. CvdB and ACD are supported in part by PPARC.

7. Referenes

[1] A. Riess et al., astro-ph/0402512
[2] C. Wetterich, Nucl. Phys. **B302**, 668 (1988)
[3] P.J.E. Peebles and B. Ratra, ApJ **325**, L17 (1988)
[4] P.J.E. Peebles and B. Ratra, Rev.Mod.Phys.**75**, 559 (2003),
[5] V. Sahni, astro-ph/0403324
[6] D. Langlois, Prog.Theor.Phys.Suppl. **148**, 181 (2003)
[7] Ph. Brax and C. van de Bruck, Class.Quant.Grav. **20**, R201 (2003)
[8] F. Quevedo, Class.Quant.Grav.**19**, 5721 (2002)
[9] R. Maartens, gr-qc/0312059
[10] Ph. Brax, C. van de Bruck and A.-C. Davis, hep-th/0404011
[11] Ph. Brax, C. van de Bruck, A.-C. Davis and C.S. Rhodes, *Dark Energy from Extra Dimensions*, in: "*On the nature of dark energy*", eds. Ph. Brax, J. Martin and J.-P. Uzan, Frontier Group, p. 203
[12] J.M. Cline and J. Vinet, Phys.Rev.D **68**, 025015 (2003)
[13] M. Gasperini, F. Piazza and G. Veneziano, Phys.Rev.D **65**, 023508 (2002)
[14] K. Freese and M. Lewis, Phys.Lett.B **540**, 1 (2002)
[15] S.M. Carrol, M. Trodden, M.S. Turner, astro-ph/0306438
[16] M. Peloso, E. Poppitz, Phys.Rev.D **68**, 125009 (2003)
[17] C.S. Rhodes, C. van de Bruck, Ph. Brax and A.-C. Davis, Phys.Rev.D **68**, 083511 (2003)
[18] K. Koyama, Phys.Rev.Lett.**91**, 221301 (2003)
[19] C. Wetterich, Astronomy and Astrophysics **301**, 321 (1995)
[20] L. Amendola, Phys.Rev.D **62**, 043511 (2000)

[21] G. F. Farrar and P.J.E. Peebles, ApJ **605**, 1 (2004)
[22] M. B. Hoffman, astro-ph/0307350
[23] J. Khoury and A. Weltman, Phys.Rev.D **69**, 044026 (2004)
[24] Ph. Brax, C. van de Bruck, A.-C. Davis, J. Khoury and A. Weltman, astro-ph/0408415
[25] B. Bertotti, L. Iess and P. Tortora, Nature **425**, 374
[26] C. M. Will, Living Rev. Rel. **4**, 4 (2001)
[27] Ph. Brax and A.-C. Davis, Phys.Lett.B**497**, 289 (2001)
[28] Ph. Brax and A.-C. Davis, JHEP **0105** 007 (2001)
[29] Ph. Brax, C. van de Bruck, A.-C. Davis and C.S. Rhodes, Phys.Rev.D**67**, 023512 (2003)
[30] L. Randall and R. Sundrum, Phys.Rev.Lett. **83**, 3370 (1999)
[31] P. J. Steinhardt, L. Wang and I. Zlatev, Phys.Rev.D **59**, 123504 (1999)
[32] B. Ratra and P.J.E. Peebles, Phys.Rev.D **37**, 3406 (1988)